

## FOR ANIMAL USE ONLY

### DISULFOX L.A.

Reg. No. G3212 (Act 36/1947)

Namibia Reg. No. V00/17.1.7/649 NSQ

### INDICATIONS

**Disulfox L.A.** is for the treatment of foot rot, navel ill, joint ill and pneumonia in stock, coccidiosis and bacterial scours in calves and lambs and strangles in horses.

### COMPOSITION

Contains: Sodium sulphadimethoxine 40 % *m/v*

### CAUTION

### STORAGE

Store in a cool dark place.

### WARNINGS

- **Withdrawal period:** Do not slaughter animals for human consumption within 7 days of the last treatment.
- Milk from treated cows must not be used for human consumption within 48 hours of last treatment.
- Dispose of empty containers according to local waste disposal regulations and do not reuse for any other purpose.
- KEEP OUT OF REACH OF CHILDREN, UNINFORMED PERSONS AND ANIMALS.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

**N.B.** Treated animals must have free access to ample drinking water during treatment and for at least two days after the last treatment.

### DIRECTIONS FOR USE - USE ONLY AS DIRECTED

### ADMINISTRATION

Administer by subcutaneous or intravenous routes.

Massage the injection site to maximise absorption and to minimise swelling.

When using the intravenous route, give the injection very slowly.

Discontinue for a while at the first sight of discomfort. Continue only when all seems normal.

### DOSAGE

An initial dosage double the maintenance dosage is recommended.

Do not treat for more than 4 days.

Species	Initial Dose	Maintenance Dose
Cattle, horses, sheep, goats and pigs	1 to 2 ml/10 kg body mass	0,5 to 1 ml/10 kg body mass

**N.B.** Consult your veterinarian immediately if any uncertainty arises as to the diagnosis or dosage or if little or no improvement is affected by the treatment.

### INDICATIONS PER TARGET SPECIES

CATTLE, SHEEP AND GOATS	HORSES	PIGS	CALVES AND LAMBS
Foot rot, pneumonia, navel ill & joint ill	Pneumonia, strangles, navel ill & joint ill	Pneumonia & joint ill	Coccidiosis & bacterial scours.

**PRESENTATION**

100 ml and 500 ml glass vial.

**REGISTRATION HOLDER**

Intervet South Africa (Pty) Ltd.

20 Spartan Road

Spartan

1619, RSA

Tel: +27 (0) 11 923 9300

Fax: +27 (0) 11 392 3158

[www.msd-animal-health.co.za](http://www.msd-animal-health.co.za)

**MANUFACTURER**

Burchem Research (Pty) Ltd.

54 Electron Avenue, Isando

1619, South Africa

**DATE OF PUBLICATION OF THIS PACKAGE INSERT**

27 February 2003