

FOR ANIMAL USE ONLY

2 *



PANACUR® 4%

Reg. No. G0169 (Act 36/1947)

Namibia Reg. No. V05/18.1.1/452 NSQ

INDICATIONS

Panacur® 4% is a roundworm remedy for pigs and cattle as well as a roundworm and milk tapeworm remedy for sheep and goats.

Targets: nodular worm, brown stomach worm and wireworm.

COMPOSITION

Contains: Fenbendazole 4,0 % *m/m*

* Benzimidazole

STORAGE

Store in a cool place.

WARNINGS

- **Withdrawal period:** Do not slaughter animals for human consumption within 7 days of last treatment.
- Do not mix with any unspecified substance.
- Sheep must be inoculated against pulpy kidney (enterotoxaemia).
- Dispose of empty containers according to local waste disposal regulations and do not reuse for any other purpose.
- KEEP OUT OF REACH OF CHILDREN, UNINFORMED PERSONS AND ANIMALS.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

DIRECTIONS FOR USE - USE ONLY AS DIRECTED

The **Panacur® 4%** powder is incorporated directly into the feed at the recommended dosage rate as per the dosage instructions.

DOSAGE AND ADMINISTRATION

Cattle, Pigs, Sheep and Goats – Roundworm

5 mg active ingredient/kg body mass i.e. 1,25 g **Panacur® 4%** powder/10 kg body mass.

Sheep and Goats – Milk tapeworm

10 mg active ingredient/kg body mass i.e. 2,50 g **Panacur® 4%** powder/10 kg body mass.

DOSAGE TABLE

Body mass (kg)	Double the dose to treat milk tapeworm infestation in sheep/goats
	Dose (per gram Panacur® 4% powder) cattle, pigs, sheep and goats
10	1,25
20	2,50
50	6,25
100	12,50
200	25,00
500	62,50

Panacur® 4% can be safely used in animals before and after mating, pregnant animals, young, old and debilitated animals.

Panacur® 4% kills worm eggs present in the animal at dosing.

EFFICACY

Pigs: 5 mg/kg body mass

<i>Ascaris suum</i>	*
---------------------	---

Cattle: 5 mg/kg body mass

Roundworms	Immature	Adult
Wireworm	*	*
Brown stomach worm	*	*
Nodular worm	*	*
Hookworm	*	*
Cattle bankrupt worm	*	*

Sheep and goats: 5 mg/kg body mass

Roundworms	Immature	Adult
Wireworm •	*	*
Nodular worm •	*	*
Hookworm	*	*
Bankrupt worm	*	*
Long-necked bankrupt worm	*	*
White bankrupt worm	*	*
Large-mouthed bowel worm	*	*
Brown stomach worm •	*	*
Lungworm	*	*

Explanation:

- * Controls (≥ 90 % effective)
- Except resistant strains

Sheep and goats: 10 mg/kg body mass

Milk tapeworm (<i>Moniezia expansa</i>)	CLASS 1
--	---------

CLASS 1: 100 % effective in at least 80 % of the treated flock.

The list contains the more important endoparasite species. This anthelmintic may also be effective against other species. For more information consult your veterinarian.

PRESENTATION

1 kg powder packed in a white plastic container with a white cap.

REGISTRATION HOLDER

Intervet South Africa (Pty) Ltd.
20 Spartan Road, Spartan
1619, RSA
Tel: +27 (0) 11 923 9300
Fax: +27 (0) 11 392 3158
www.msd-animal-health.co.za

MANUFACTURER

Burchem Research (Pty) Ltd.
54 Electron Rd
Isando

DATE OF PUBLICATION OF THIS PACKAGE INSERT

28 March 2000