

FOR ANIMAL USE ONLY



PANACUR® BS

Reg. No. G1481 (Act 36/1947)

Namibia Reg. No. V03/18.1.1/655 NSO

INDICATION

- A Roundworm and lungworm remedy for cattle.
- A Roundworm, lungworm and milk tapeworm remedy for sheep and goats.
- A Roundworm remedy for horses and ostriches.

CAUTION

COMPOSITION

Contains: Fenbendazole 5 % *m/v*

(*Benzimidazole)

STORAGE

Store in a cool, dry place.

WARNINGS

- **Withdrawal periods:**
 - **Cattle:** Meat – 7 days
Milk – 3 days
 - **Sheep & Goats:** Meat – 14 days
Milk – 4 days
 - **Ostriches:** Meat – 7 Days
- Do not mix or dilute with any unspecified substance.
- Sheep must be inoculated against pulpy kidney (enterotoxemia) before dosing.
- Keep away from foodstuffs.
- Destroy empty containers according to local waste disposal regulations and do not reuse for any other purpose.
- KEEP OUT OF REACH OF CHILDREN, UNINFORMED PERSONS AND ANIMALS.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

DIRECTIONS FOR USE - USE ONLY AS DIRECTED

- Shake well before use.
- Ready-to-use.

1. CATTLE & HORSES

a) DOSAGE RATE

Cattle: Dose **5 mg/kg** body mass i.e. **1 ml/10 kg** body mass

Horses: Dose **10 mg/kg** body mass i.e. **2 ml/10 kg** body mass

b) DOSAGE TABLE

| BODY MASS | DOSE (mℓ) | | DOSAGES IN PACK (200 mℓ) | |
|-----------|-----------|--------|-----------------------------|--------|
| | CATTLE | HORSES | CATTLE | HORSES |
| 50 | 5 | 10 | 40 | 20 |
| 100 | 10 | 20 | 20 | 10 |
| 150 | 15 | 30 | 13 | 6 |
| 200 | 20 | 40 | 10 | 5 |
| 250 | 25 | 50 | 8 | 4 |
| 300 | 30 | 60 | 7 | 3 |

| BODY MASS | DOSE (mℓ) | | DOSAGES IN PACK (1 ℓ) | |
|-----------|-----------|--------|--------------------------|--------|
| | CATTLE | HORSES | CATTLE | HORSES |
| 50 | 5 | 10 | 200 | 100 |
| 100 | 10 | 20 | 100 | 50 |
| 150 | 15 | 30 | 66 | 33 |
| 200 | 20 | 40 | 50 | 25 |
| 250 | 25 | 50 | 40 | 20 |
| 300 | 30 | 60 | 33 | 16 |

| BODY MASS | DOSE (mℓ) | | DOSAGES IN PACK (5 ℓ) | |
|-----------|-----------|--------|--------------------------|--------|
| | CATTLE | HORSES | CATTLE | HORSES |
| 50 | 5 | 10 | 1 000 | 500 |
| 100 | 10 | 20 | 500 | 250 |
| 150 | 15 | 30 | 333 | 166 |
| 200 | 20 | 40 | 250 | 125 |
| 250 | 25 | 50 | 200 | 100 |
| 300 | 30 | 60 | 166 | 83 |

| BODY MASS | DOSE (mℓ) | | DOSAGES IN PACK (10 ℓ) | |
|-----------|-----------|--------|---------------------------|--------|
| | CATTLE | HORSES | CATTLE | HORSES |
| 50 | 5 | 10 | 4 000 | 2 000 |
| 100 | 10 | 20 | 2 000 | 1 000 |
| 150 | 15 | 30 | 1 333 | 666 |
| 200 | 20 | 40 | 1 000 | 500 |
| 250 | 25 | 50 | 800 | 400 |
| 300 | 30 | 60 | 666 | 333 |

Panacur® BS can be safely used in all classes of stock and horses.

c) **EFFICACY In Cattle** at 5 mg fenbendazole/kg body mass.

| ROUNDWORM SPECIES | IMMATURE | ADULT |
|--|-----------------|--------------|
| Wireworm (<i>Haemonchus placei</i>) | * | * |
| Brown stomach worm (<i>Ostertagia ostertagi</i>) | * | * |
| Cattle bankrupt worm (<i>Cooperia</i> spp.) | * | * |
| Nodular worm (<i>Oesophagostomum radiatum</i>) | * | * |
| Hookworm (<i>Bunostomum plebotomum</i>) | * | * |
| Lungworm (<i>Dictyocaulus viviparus</i>) | * | * |

Explanation

* Controls (≥ 90 % effective)

** Aids in the control (60 to 89 % effective)

d) **EFFICACY In Horses** at 10 mg fenbendazole/kg body mass.

Panacur® BS at the recommended dose is highly effective against the following roundworm species:

- Bloodworm (large strongyles)
- Redworm (small strongyles)
- Whipworms (pinworm)
- Ascarids
- Stomach worm (*Habronema majus*)

The list contains the more important endoparasite species. This anthelmintic may also be effective against other species. For more information consult your veterinarian.

2. SHEEP & GOATS

a) **DOSAGE RATE**

- For round- and lungworm dose at **5 mg/kg** body mass i.e. **1 ml/10 kg** body mass.
- For round-, lung- and milk tapeworm dose at **10 mg/kg** body mass i.e. **1 ml/5 kg** body mass.
- Double the dosage and halve the number of doses per pack when treating for milk tapeworm.

b) DOSAGE TABLE

| BODY MASS (kg) | DOSE (mℓ) | DOSES IN PACK (200 mℓ) |
|-----------------------|------------------|-----------------------------------|
| 5 | ½ | 400 |
| 10 | 1 | 200 |
| 15 | 1½ | 133 |
| 20 | 2 | 100 |
| 30 | 3 | 66 |
| 40 | 4 | 50 |
| 50 | 5 | 40 |
| 60 | 6 | 33 |

| BODY MASS (kg) | DOSE (mℓ) | DOSES IN PACK (1 ℓ) |
|-----------------------|------------------|--------------------------------|
| 5 | ½ | 2 000 |
| 10 | 1 | 1 000 |
| 15 | 1½ | 666 |
| 20 | 2 | 500 |
| 30 | 3 | 333 |
| 40 | 4 | 250 |
| 50 | 5 | 200 |
| 60 | 6 | 166 |

| BODY MASS (kg) | DOSE (mℓ) | DOSES IN PACK (5 ℓ) |
|-----------------------|------------------|--------------------------------|
| 5 | ½ | 10 000 |
| 10 | 1 | 5 000 |
| 15 | 1½ | 3 333 |
| 20 | 2 | 2 500 |
| 30 | 3 | 1 666 |
| 40 | 4 | 1 250 |
| 50 | 5 | 1 000 |
| 60 | 6 | 833 |

| BODY MASS (kg) | DOSE (mℓ) | DOSES IN PACK (10 ℓ) |
|-----------------------|------------------|---------------------------------|
| 5 | ½ | 20 000 |
| 10 | 1 | 10 000 |
| 15 | 1½ | 6 666 |
| 20 | 2 | 5 000 |
| 30 | 3 | 3 333 |
| 40 | 4 | 2 500 |
| 50 | 5 | 2 000 |
| 60 | 6 | 1 666 |

Panacur® BS can be used safely in rams and ewes before and after mating, pregnant ewes, young lambs, old and debilitated animals. Safe even at 50 times the recommended dose. Does not stain wool or mohair. Kills worm eggs present in the animal at dosing.

EFFICACY IN SHEEP AND GOATS

Roundworms at 5 mg fenbendazole/kg body mass.

| ROUNDWORM SPECIES | IMMATURE | ADULT |
|---|----------|-------|
| Wireworm† (<i>Haemonchus contortus</i>) | * | * |
| Bankrupt worm (<i>Trichostrongylus colubriformis</i>) | * | * |
| Longnecked bankrupt worm (<i>Nematodirus spathiger</i>) | * | * |
| Brown stomach worm† (<i>Teladorsagia circumcincta</i>) | * | * |
| Nodular worm (<i>Oesophagostomum columbianum</i>) | * | * |
| Hookworm† (<i>Gaigeria pachyscelis</i>) | * | * |
| Large-mouthed bowelworm (<i>Chabertia ovina</i>) | * | * |
| White bankrupt worm (<i>Strongyloides papillosus</i>) | * | * |
| Lungworm (<i>Dictyocaulus filaria</i>) | * | * |

Explanation

- * Controls (≥ 90 % effective)
- ** Aids in the control (60 to 89 % effective)
- † Except resistant strains

Tapeworms at 10 mg fenbendazole/kg body mass.

| TAPEWORM SPECIES | EFFICACY CLASS |
|--|----------------|
| Milk Tapeworm (<i>Moniezia expansa</i>) | 1 |

Explanation

- CLASS 1: 100 % effective in at least 80 % of the treated flock
- CLASS 2: 100 % effective in at least 60 % of the treated flock
- CLASS 3: 100 % effective in at least 50 % of the treated flock
- CLASS 4: Ineffective

The list contains the more important endoparasite species. This anthelmintic may also be effective against other species. For more information consult your veterinarian.

3. OSTRICHES

a) DOSAGE RATE

- Dose 1,5 mℓ/5 kg body mass.

b) DOSAGE TABLE

| BODY MASS (kg) | DOSE (mℓ) | DOSES IN PACK (200 mℓ) |
|----------------|-----------|---------------------------|
| 5 | 1½ | 133 |
| 10 | 3 | 66 |
| 20 | 6 | 33 |
| 30 | 9 | 22 |
| 40 | 12 | 16 |
| 50 | 15 | 13 |

| BODY MASS (kg) | DOSE (mℓ) | DOSES IN PACK (1 ℓ) |
|----------------|-----------|------------------------|
| 5 | 1½ | 666 |
| 10 | 3 | 333 |
| 20 | 6 | 166 |
| 30 | 9 | 111 |
| 40 | 12 | 83 |
| 50 | 15 | 66 |

| BODY MASS (kg) | DOSE (mℓ) | DOSES IN PACK (5 ℓ) |
|----------------|-----------|------------------------|
| 5 | 1½ | 3 333 |
| 10 | 3 | 1 666 |
| 20 | 6 | 833 |
| 30 | 9 | 555 |
| 40 | 12 | 416 |
| 50 | 15 | 333 |

| BODY MASS (kg) | DOSE (mℓ) | DOSES IN PACK (10 ℓ) |
|----------------|-----------|-------------------------|
| 5 | 1½ | 6 666 |
| 10 | 3 | 3 333 |
| 20 | 6 | 1 666 |
| 30 | 9 | 1 111 |
| 40 | 12 | 833 |
| 50 | 15 | 666 |

c) EFFICACY in Ostriches

| OSTRICH ROUNDWORM | |
|-----------------------------------|---|
| <i>Libyostrongylus douglassii</i> | * |

* Controls (≥ 90 % effective)

Panacur® BS is safe for mature ostriches and ostrich chicks. The list contains the more important endoparasite species. This anthelmintic may also be effective against other species. For more information consult your veterinarian.

PRESENTATION

200 ml, 1 l, 5 l and 10 l unfluorinated plastic bottles, sealed with an EXPE lined cap.

REGISTRATION HOLDER

Intervet South Africa (Pty) Ltd.
20 Spartan Road, Spartan
1619, RSA
Tel: +27 (0) 11 923 9300
Fax: +27 (0) 11 392 3158
www.msd-animal-health.co.za

MANUFACTURER

Burchem Research (Pty) Ltd.
54 Electron Rd, Kempton Park
1619, South Africa

DATE OF PUBLICATION OF THIS PACKAGE INSERT

5 February 2018

| |
|---|
| Zimbabwe Reg.No. 94/80.12.10/9360 Pharmaceutical Classification: 801210 Distribution category: VMGD |
|---|

SLEGS VIR DIEREGERBUIK



PANACUR® BS

Reg. Nr. G1481 (Wet 36/1947)

Namibië Reg. Nr. V03/18.1.1/655 NSQ

INDIKASIE

- 'n Ronde- en Longwurmmiddel vir beeste.
- 'n Ronde-, Long- en Melklintwurmmiddel vir skape en bokke.
- 'n Rondewurmmiddel vir perde en volstruise.

VERSIGTIG

SAMESTELLING

Bevat: Fenbendasool 5 % *m/v*

(*Bensimidiasool)

BERGING

Bewaar in a koel, droë plek.

WAARSKUWINGS

- **Ontrekkingsperiodes:**
 - **Beeste:** Vleis – 7 dae
Melk – 3 dae
 - **Skape & Bokke:** Vleis – 14 dae
Melk – 4 dae
 - **Volstruise:** Vleis – 7 dae
- Moenie met enige ongespesifiseerde middel gemeng of verdun word nie.
- Skape moet voor dosering teen bloednier (enterotoksemie) ingeënt word.
- Hou weg van voedsel.
- Doen weg met enige leë houers volgens plaaslike afvalbestuursregulasies en moenie vir enige ander doeleinde hergebruik nie.
- HOU BUIE DIE BEREIK VAN KINDERS, ONINGELIGTE PERSONE EN DIERE.
- Alhoewel hierdie middel breedvoerig onder 'n wye verskeidenheid toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.

GEbruiksaanwysings - GEbruik SLEGS SOOS AANGEDUI

- Skud goed voor gebruik.
- Gereed-vir-gebruik.

1. BEESTE & PERDE

a) DOSIS

Beeste: Doseer **5 mg/kg** liggaamsmassa, m.a.w. **1 ml/10 kg** liggaamsmassa.

Perde: Doseer **10 mg/kg** liggaamsmassa, m.a.w. **2 ml/10 kg** liggaamsmassa.

b) Doseringstabel

| LIGGAAMSMASSA | DOSIS (mℓ) | | DOSIS IN VERPAKKING (200 mℓ) | |
|---------------|------------|-------|---------------------------------|-------|
| | BEESTE | PERDE | BEESTE | PERDE |
| 50 | 5 | 10 | 40 | 20 |
| 100 | 10 | 20 | 20 | 10 |
| 150 | 15 | 30 | 13 | 6 |
| 200 | 20 | 40 | 10 | 5 |
| 250 | 25 | 50 | 8 | 4 |
| 300 | 30 | 60 | 7 | 3 |

| LIGGAAMSMASSA | DOSIS (mℓ) | | DOSIS IN VERPAKKING (1 ℓ) | |
|---------------|------------|-------|------------------------------|-------|
| | BEESTE | PERDE | BEESTE | PERDE |
| 50 | 5 | 10 | 200 | 100 |
| 100 | 10 | 20 | 100 | 50 |
| 150 | 15 | 30 | 66 | 33 |
| 200 | 20 | 40 | 50 | 25 |
| 250 | 25 | 50 | 40 | 20 |
| 300 | 30 | 60 | 33 | 16 |

| LIGGAAMSMASSA | DOSIS (mℓ) | | DOSIS IN VERPAKKING (5 ℓ) | |
|---------------|------------|-------|------------------------------|-------|
| | BEESTE | PERDE | BEESTE | PERDE |
| 50 | 5 | 10 | 1 000 | 500 |
| 100 | 10 | 20 | 500 | 250 |
| 150 | 15 | 30 | 333 | 166 |
| 200 | 20 | 40 | 250 | 125 |
| 250 | 25 | 50 | 200 | 100 |
| 300 | 30 | 60 | 166 | 83 |

| LIGGAAMSMASSA | DOSIS (mℓ) | | DOSIS IN VERPAKKING (10 ℓ) | |
|---------------|------------|-------|-------------------------------|-------|
| | BEESTE | PERDE | BEESTE | PERDE |
| 50 | 5 | 10 | 4 000 | 2 000 |
| 100 | 10 | 20 | 2 000 | 1 000 |
| 150 | 15 | 30 | 1 333 | 666 |
| 200 | 20 | 40 | 1 000 | 500 |
| 250 | 25 | 50 | 800 | 400 |
| 300 | 30 | 60 | 666 | 333 |

Panacur® BS kan met veiligheid by alle klasse van vee en perde gebruik word.

c) **DOELTREFFENDHEID in Beeste** teen 5 mg fenbendasool/kg liggaamsmassa

| RONDEWURMSPECIES | ONVOLWASSE | VOLWASSE |
|--|------------|----------|
| Haarwurm (<i>Haemonchus placei</i>) | * | * |
| Bruinmaagwurm (<i>Ostertagia ostertagi</i>) | * | * |
| Beesbankrotwurm (<i>Cooperia</i> spp.) | * | * |
| Knoppieswurm (<i>Oesophagostomum radiatum</i>) | * | * |
| Haakwurm (<i>Bunostomum plebotomum</i>) | * | * |
| Longwurm (<i>Dictyocaulus viviparus</i>) | * | * |

Verduideliking

* Beheer (≥ 90 % effektief)

** Help in die beheer (60 tot 89 % effektief)

d) **DOELTREFFENDHEID In Perde** teen 10 mg fenbendasool/kg liggaamsmassa

Panacur® BS teen die aanbevole dosis is hoogs effektief teen die volgende rondewurmspesies:

- Bloedwurms (groot strongiele)
- Rooiwurm (klein strongiele)
- Sambokwurm
- Spoelwurm
- Maagwurm (*Habronema majus*)

Hierdie lys bevat die belangrikste endoparasitiese spesies. Hierdie wurmmiddel kan ook effektief wees teen ander spesies. Vir meer inligting raadpleeg u veearts.

1. **SKAPE & BOKKE**

a) **DOSIS**

- Vir ronde- en longwurm doseer teen **5 mg/kg** liggaamsmassa m.a.w. **1 ml/10 kg** liggaamsmassa.
- Vir ronde-, long- en melklintwurm doseer teen **10 mg/kg** liggaamsmassa m.a.w. **1 ml/5 kg** liggaamsmassa.
- Verdubbel die dosisse en halveer die aantal dosisse per verpakking vir behandeling van melklintwurm.

b) DOSERINGSTABEL

| LIGGAAMSMASSA (kg) | DOSIS (mℓ) | DOSIS IN VERPAKKING (200 mℓ) |
|--------------------|------------|------------------------------|
| 5 | ½ | 400 |
| 10 | 1 | 200 |
| 15 | 1½ | 133 |
| 20 | 2 | 100 |
| 30 | 3 | 66 |
| 40 | 4 | 50 |
| 50 | 5 | 40 |
| 60 | 6 | 33 |

| LIGGAAMSMASSA (kg) | DOSIS (mℓ) | DOSIS IN VERPAKKING (1 ℓ) |
|--------------------|------------|---------------------------|
| 5 | ½ | 2 000 |
| 10 | 1 | 1 000 |
| 15 | 1½ | 666 |
| 20 | 2 | 500 |
| 30 | 3 | 333 |
| 40 | 4 | 250 |
| 50 | 5 | 200 |
| 60 | 6 | 166 |

| LIGGAAMSMASSA (kg) | DOSIS (mℓ) | DOSIS IN VERPAKKING (5 ℓ) |
|--------------------|------------|---------------------------|
| 5 | ½ | 10 000 |
| 10 | 1 | 5 000 |
| 15 | 1½ | 3 333 |
| 20 | 2 | 2 500 |
| 30 | 3 | 1 666 |
| 40 | 4 | 1 250 |
| 50 | 5 | 1 000 |
| 60 | 6 | 833 |

| LIGGAAMSMASSA (kg) | DOSIS (mℓ) | DOSIS IN VERPAKKING (10 ℓ) |
|--------------------|------------|----------------------------|
| 5 | ½ | 20 000 |
| 10 | 1 | 10 000 |
| 15 | 1½ | 6 666 |
| 20 | 2 | 5 000 |
| 30 | 3 | 3 333 |
| 40 | 4 | 2 500 |
| 50 | 5 | 2 000 |
| 60 | 6 | 1 666 |

Panacur® BS kan met veiligheid by ramme en ooie voor en na paring, dragtige ooie, jong lammers, oud en verswakte diere gebruik word. Veilig selfs teen 50 maal die aanbevole dosis. Sal nie wol of sybokhaar vlek nie. Dood wurmeiers wat tydens dosering in die dier aanwesig is.

c) DOELTREFFENDHEID In Skape en Bokke

RONDEWURMS teen 5 mg fenbendasool/kg liggaamsmassa

| RONDEWURMSPEISIE | ONVOLWASSE | VOLWASSE |
|---|------------|----------|
| Haarwurm† (<i>Haemonchus contortus</i>) | * | * |
| Bankrotwurm (<i>Trichostrongylus colubriformis</i>) | * | * |
| Langnekbankrotwurm (<i>Nematodirus spathiger</i>) | * | * |
| Bruinmaagwurm† (<i>Teladorsagia circumcincta</i>) | * | * |
| Knoppieswurm (<i>Oesophagostomum columbianum</i>) | * | * |
| Haakwurm† (<i>Gaigeria pachyscelis</i>) | * | * |
| Grootbekwurm (<i>Chabertia ovina</i>) | * | * |
| Witbankrotwurm (<i>Strongyloides papillosus</i>) | * | * |
| Longwurm (<i>Dictyocaulus filaria</i>) | * | * |

Verduideliking

- * Beheer (≥ 90 % effektief)
- ** Help in die beheer (60 tot 89 % effektief)
- † Behalwe weerstandbiedende stamme

LINTWURMS teen 10 mg fenbendasool/kg liggaamsmassa

| LINTWURMSPEISIE | EFFEKTIWITEIT KLAS 1 |
|---|----------------------|
| Melklintwurm (<i>Moniezia expansa</i>) | 1 |

Verduideliking

- KLAS 1: 100 % effektief in minstens 80 % van die behandelde trop
- KLAS 2: 100 % effektief in minstens 60 % van die behandelde trop
- KLAS 3: 100 % effektief in minstens 50 % van die behandelde trop
- KLAS 4: Oneffektief

Hierdie lys bevat die belangrikste endoparasitiese spesies. Hierdie wurmmiddel kan ook effektief wees teen ander spesies. Vir meer inligting raadpleeg jou veearts.

2. VOLSTRUISE

a) DOSIS

- Doseer **1,5 ml/5 kg** liggaamsmassa.

b) DOSERINGSTABEL

| LIGGAAMSMASSA (kg) | DOSIS (mℓ) | DOSIS IN VERPAKKING (200 mℓ) |
|--------------------|------------|------------------------------|
| 5 | 1½ | 133 |
| 10 | 3 | 66 |
| 20 | 6 | 33 |
| 30 | 9 | 22 |
| 40 | 12 | 16 |
| 50 | 15 | 13 |

| LIGGAAMSMASSA (kg) | DOSIS (mℓ) | DOSIS IN VERPAKKING (1 ℓ) |
|--------------------|------------|---------------------------|
| 5 | 1½ | 666 |
| 10 | 3 | 333 |
| 20 | 6 | 166 |
| 30 | 9 | 111 |
| 40 | 12 | 83 |
| 50 | 15 | 66 |

| LIGGAAMSMASSA (kg) | DOSIS (mℓ) | DOSIS IN VERPAKKING (5 ℓ) |
|--------------------|------------|---------------------------|
| 5 | 1½ | 3 333 |
| 10 | 3 | 1 666 |
| 20 | 6 | 833 |
| 30 | 9 | 555 |
| 40 | 12 | 416 |
| 50 | 15 | 333 |

| LIGGAAMSMASSA (kg) | DOSIS (mℓ) | DOSIS IN VERPAKKING (10 ℓ) |
|--------------------|------------|----------------------------|
| 5 | 1½ | 6 666 |
| 10 | 3 | 3 333 |
| 20 | 6 | 1 666 |
| 30 | 9 | 1 111 |
| 40 | 12 | 833 |
| 50 | 15 | 666 |

c) DOELTREFFENDHEID in Volstruise

| VOLSTRUIS RONDEWURM | |
|-----------------------------------|---|
| <i>Libyostrongylus douglassii</i> | * |

* Beheer (≥ 90 % effektief)

Panacur® BS is veilig vir volwasse volstruise en volstruiskuikens. Hierdie wurmmiddel kan ook effektief wees teen ander spesies. Vir meer inligting raadpleeg jou veearts.

AANBIEDING

200 ml, 1 l, 5 l en 10 l ongefloreerde plastiekbottels, verseël met 'n EXPE-gevoerde doppie.

REGISTRASIEHOUER

Intervet South Africa (Pty) Ltd.

Spartanweg 20, Spartan

1619, RSA

Tel: +27 (0) 11 923 9300

Faks: +27 (0) 11 392 3158

www.msd-animal-health.co.za

VERVAARDIGER

Burchem Research (Pty) Ltd.

54 Electron Rd, Kempton Park

1619, South Africa

DATUM VAN PUBLIKASIE VAN HIERDIE VOUBILJET

5 Februarie 2018