

FOR ANIMAL USE ONLY

2*

PANACUR® PASTE

Reg. No. G3374 (Act 36/1947)

Namibia Reg. No. V04/18.1.1/248 NS0

INDICATIONS

For the treatment of immature and mature stages of roundworms: bloodworms (large strongyles), redworms (small strongyles), ascarids, oxyurids and strongyloides of horses.

COMPOSITION

Each 1 g of paste contains: Fenbendazole 187,5 mg

* Benzimidazole

STORAGE

Store at room temperature between 20 °C and 25 °C.

WARNINGS

- Dispose of all empty syringes according to local waste disposal regulations and do not reuse for any other purpose.
- KEEP OUT OF REACH OF CHILDREN, UNINFORMED PERSONS AND ANIMALS.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

DIRECTIONS FOR USE - USE ONLY AS DIRECTED

- The syringe is graduated with 6 markings. Each marking corresponds to a dosage for 100 kg body weight.
- Before administration, the syringe should be adjusted to the calculated dosage by setting the ring on the plunger.
- Ensure that the horse does not have food in its mouth.
- Insert the syringe in the gap between the front and back teeth and deliver paste onto the back of the tongue.
- Immediately raise the horse's head for a few seconds after dosing.

DOSAGE

- The recommended dosage is 7,5 mg fenbendazole/kg body weight. One syringe contains sufficient paste to treat one 600 kg horse.
- Diarrhoea caused by *Strongyloides westeri* in suckling foals should be treated with a dose of 50 mg fenbendazole/kg body weight corresponding to 24 g (contents of 1 syringe) for 90 kg body weight.

EFFICACY

Roundworms		Stage/s	Controls* /Aids in the control**	Dosage Rate
Small strongyles including:	<i>Craterostomum</i> spp., <i>Cylicocyclus</i> spp., <i>Cylicodontophorus</i> spp., <i>Cylicostephanus</i> spp., <i>Cyathostomum</i> spp., <i>Gyalocephalus</i> spp., <i>Poteriostomum</i> spp.	} Adult	*	7,5 mg/kg/day for 1 day
Large strongyles including:	<i>Oesophagodontus</i> spp.	Adult	*	7,5 mg/kg/day for 1 day
	<i>Strongylus edentatus</i>	Larval Stages	*	7,5 mg/kg/day for 5 days
		Adult	*	7,5 mg/kg/day for 1 day
	<i>S. equinus</i>	Adult	*	7,5 mg/kg/day for 1 day
	<i>S. vulgaris</i>	Larval Stages in arteries	**	7,5 mg/kg/day for 5 days
		Adult	*	7,5 mg/kg/day for 1 day
	<i>S. westeri</i>	Adult	*	50 mg/kg/day for 1 day
<i>Triodontophorus</i>	Adult	*	7,5 mg/kg/day for 1 day	
Pinworms (<i>Oxyuris equi</i>)	4 th Stage larvae		*	7,5 mg/kg/day for 1 day
	Adult		*	7,5 mg/kg/day for 1 day
Ascarids (<i>Parascaris equorum</i>)	Adult		*	7,5 mg/kg/day for 1 day
Small strongyles	Immature (mucosal stages)		*	7,5 mg/kg/day for 5 days
<i>Habronema</i> spp. and <i>Draschia</i> spp.	Adult		*	7,5 mg/kg/day for 5 days
<i>Probstmayria viviparus</i>	Adult		*	7,4 mg/kg/day for 1 day

* Controls (≥ 90 % effective)

** Aids in the control (60 to 89 % effective)

Panacur® Paste can be used in adult horses, pregnant and lactating mares, foals and stallions.

PRESENTATION

24 g syringes.

REGISTRATION HOLDER

Intervet South Africa (Pty) Ltd.
20 Spartan Road, Spartan
1619, RSA
Tel: +27 (0) 11 923 9300
Fax: +27 (0) 11 392 3158
www.msd-animal-health.co.za

MANUFACTURER

Intervet Productions S.A.
Rue de Lyons
Igoville, France

DATE OF PUBLICATION OF THIS PACKAGE INSERT

1 December 2003