

FOR ANIMAL USE ONLY

TRIVETRIN® INJECTION

Reg. No. G1742 (Act 36/1947)

Namibia Reg. No. V03/17.1.7/516 NS0

INDICATIONS

Trivetrim® Injection is effective against a wide range of gram-positive and gram-negative organisms including *Staphylococcus* (including most penicillinase-producing strains), *Streptococcus*, *Salmonella*, *E. coli*, *Haemophilus*, *Proteus*, *Pasteurella*, *Klebsiella* and *Enterobacter*.

Trivetrim® Injection is indicated for the treatment of bacterial infections of the respiratory tract, urinary tract, genital tract and alimentary tract. **Trivetrim® Injection** is also highly effective in the treatment of foot rot, severe mastitis, bacterial agalactia of sows and the treatment of wound infections and septicaemia.

COMPOSITION

Each 1 ml contains: Sulfadoxine 200 mg and Trimethoprim 40 mg.

Preservative: Diethanolamine 3 mg.

STORAGE

- Store below 25 °C in a cool, dark place.
- Do not freeze.

WARNINGS

- **Withdrawal periods -**

Tissues

Cattle: 9 days

Pig: 8 days

Sheep: 14 days

Horse: 10 days

Milk

Cattle: 3 days/72 hours

Sheep: 4,5 days/108 hours

Do not use in horses producing milk for human consumption.

- **Trivetrim® Injection** is not suitable for cats and kittens.
- Do not administer to animals with a history of sulphonamide sensitivity or showing marked liver parenchymal damage.
- Do not use **Trivetrim® Injection** in conjunction with detomidine hydrochloride.
- Do not administer subcutaneously.
- Dispose of any empty containers, needles and syringes as per local waste disposal regulations and do not reuse for any other purpose.
- KEEP OUT OF REACH OF CHILDREN, UNINFORMED PERSONS AND ANIMALS.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

DIRECTIONS FOR USE - USE ONLY AS DIRECTED.

Dosage and Administration route

Cattle, sheep and pigs: Intramuscular injection

May be given by slow intravenous injection in cases of acute infection. Consult your veterinarian.

Dogs and horses: Slow intravenous injection only.

Dose

Normal: 1 ml/16 kg body mass daily (15 mg/kg)

Severe infection: 1 ml/10 kg body mass daily (24 mg/kg)

SIDE-EFFECTS

Occasionally local swellings of a temporary nature may develop at injection sites.

KNOWN SYMPTOMS OF OVERDOSAGE

None.

PRESENTATION

100 ml amber glass vial containing a clear solution.

REGISTRATION HOLDER

Intervet South Africa (Pty) Ltd.

20 Spartan Road, Spartan

1619, RSA

Tel: +27 (0) 11 923 9300

Fax: +27 (0) 11 392 3158

www.msd-animal-health.co.za

MANUFACTURER

Vet Pharma Friesoythe GmbH

Sedelsbergerstreet 2-4

26169 Friesoythe, Germany

DATE OF PUBLICATION OF THIS PACKAGE INSERT

31 January 2018

Zimbabwe registration number: E94/80.22.20/9359

Pharmacological classification: 802220

(Sulphonamides and sulphonamide combinations)

Distribution category: PP (Vet)